

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

 Maxforce[®] Fusion
COCKROACH GEL
ACTIVE CONSTITUENT: 21.5 g/kg IMIDACLOPRID

GROUP	4A	INSECTICIDE
--------------	-----------	--------------------

**30 g - 50 g
NET**

For the control of cockroaches in the situations specified in the directions for use.

The syringe must not be sold separately. DO NOT destroy carton / blister pack and booklet while product still remains.

DIRECTIONS FOR USE

Situations	Pest	Rate	Critical Comments
Barracks, farm buildings, food processing establishments, factories, homes, hospitals, meatworks (non-product areas), offices, restaurants, schools, shops, storerooms, transport vehicles & facilities	<u>Small cockroaches</u> e.g. German Cockroach (<i>Blattella germanica</i>)	<u>Light infestations</u> 1 placement per m ² <u>Heavy infestations</u> 2 placements per m ² 1 placement = a 5 mm ball (about the size of a small pea or approx. 0.1 g) or a line about 2.5 cm long	Apply gel in out of the way locations where cockroaches occur e.g., under fridges, near water heaters, behind microwave ovens, garbage bins etc. Re-assessment of treated areas one month after initial baiting is recommended to evaluate if additional baiting is required. Bait placements should be replenished when consumed or at least every three months as long as pest populations persist.
	<u>Large cockroaches</u> e.g. American Cockroach and Australian Cockroach (<i>Periplaneta</i> spp.)	<u>Light infestations</u> 2 placements per m ² <u>Heavy infestations</u> 3 placements per m ²	<i>Periplaneta</i> spp. require more bait because of the quantity which can be consumed by a single cockroach. Reassess baiting effectiveness as above.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

DO NOT allow to contact food, livestock feed or their water. DO NOT place gel on surfaces where food is prepared or on utensils that are used for food preparation, on food packing materials or on surfaces where food packing materials are stored.

GENERAL INSTRUCTIONS

Apply this product using a suitable application device. To apply the gel, remove the cap on the nozzle and touch the tip to the surface to be treated and slowly depress plunger until sufficient gel is dispensed. Recap the dispenser after treatment is completed. Gel should be applied out of reach of children, pets and other animals. DO NOT place gel on surfaces which are regularly washed.

INSECTICIDE RESISTANCE WARNING

GROUP	4A	INSECTICIDE
--------------	-----------	--------------------

For insecticide resistance management, Maxforce Fusion Cockroach Gel is a Group 4A insecticide. Some naturally occurring insect biotypes resistant to Maxforce Fusion and other Group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Maxforce Fusion or other Group 4A insecticides are used repeatedly. The effectiveness of Maxforce Fusion on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience Pty Ltd. accepts no liability for any losses that may result from the failure of Maxforce Fusion to control resistant insects. Maxforce Fusion may be subject to specific resistance management strategies. For further information contact your local supplier or Bayer Environmental Science representative.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the gel or used syringes.

STORAGE AND DISPOSAL

Keep out of reach of children. DO NOT store in direct sunlight. Store in the closed, original carton / blister pack, in a cool dry place. Completely empty syringe in normal use. Dispose of empty syringe and carton / blister pack by wrapping in paper, placing in plastic bag and putting in garbage.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (phone Australia 13 11 26).

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which is available at www.bayeres.com.au.

Exclusion of Liability

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty. Ltd. accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

APVMA Approval No. 69488 / 113374

Maxforce® is a Registered Trademark of the Bayer Group.

Bayer CropScience Pty. Ltd.
A.B.N. 87 000 226 022
Level 1, 8 Redfern Rd,
Hawthorn East Vic. 3123
Technical enquiries: 1800 804 479
Website: www.environmentalscience.bayer.com.au

For 24 hour specialist advice in an emergency only phone 1800 033 111

Batch Number:

DOM:

Barcode

GHS Classification not required on this label

Syringe label

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

 Maxforce[®] Fusion
COCKROACH GEL
ACTIVE CONSTITUENT: 21.5 g/kg IMIDACLOPRID

GROUP	4A	INSECTICIDE
--------------	-----------	--------------------

**30 g – 50 g
NET**

For the control of cockroaches in the situations specified in the Directions for Use.

Before using product read directions on the outer pack / booklet. Sale of this syringe except in outer carton / blister pack is illegal. DO NOT destroy carton / blister pack and booklet while product still remains.

STORAGE AND DISPOSAL

Keep out of reach of children. DO NOT store in direct sunlight. Store in the closed, original carton, in a cool dry place. Completely empty syringe in normal use. Dispose of empty syringe and carton / blister pack by wrapping in paper, placing in plastic bag and putting in garbage.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone 13 11 26).

APVMA Approval No. 69488 / 113374

Maxforce[®] is a Registered Trademark of the Bayer Group.

Bayer CropScience Pty. Ltd.
A.B.N. 87 000 226 022
Level 1, 8 Redfern Rd,
Hawthorn East Vic. 3123
Technical enquiries: 1800 804 479
Website: www.environmentalscience.bayer.com.au