Product Name: APVMA Approval No:

BIOPREN 50 LIQUID MOSQUITO LARVICIDE 62020/126154



Label Name: BIOPREN 50 LIQUID MOSQUITO LARVICIDE
--

Signal Headings: READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent Statements:	50 g/L (S)-METHOPRENE
----------------------------	-----------------------

Mode of Action:			
	GROUP	7A	INSECTICIDE

Statement of Claims: For the prevention of adult mosquito emergence

Net Contents:	10L(2x5L) 5L
---------------	-----------------

Restraints:	Restraint: DO NOT apply to dry sites. Biopren 50 Liquid Mosquito Larvicide must be applied directly to water.
-------------	--

Directions for Use:	This section contains file attachment.

	Other Limitations:	Users in NSW must comply with the NSW Clean Waters Act 1970.	
--	--------------------	--	--

Withholding Periods:	WITHHOLDING PERIOD (MEAT): NO WITHHOLDING PERIOD IS REQUIRED FOR LIVESTOCK DRINKING TREATED WATER.
----------------------	--

Trade Advice:				
---------------	--	--	--	--

General Instructions:	GENERAL INSTRUCTIONS BIOPREN 50 Liquid Mosquito Larvicide releases (S)-methoprene Insect Growth Regulator into water where mosquitoes breed. BIOPREN 50 Liquid Mosquito Larvicide prevents the emergence of adult mosquitoes including common Aedes, Anopheles, Culex and Ocherotatus spp. from treated sites.
	(S)-methoprene has no effect on mosquitoes which have reached the pupal or adult stage prior to treatment.
	Treated larvae continue to develop normally to the pupal stage after which they fail to emerge. BIOPREN 50 Liquid Mosquito Larvicide has sufficient activity to be effective against existing broods when applied to any larval stage at recommended rates. Third and fourth larval stages are more susceptible to the effects of (S)-methoprene than early stages.
	For aerial application, BIOPREN 50 Liquid Mosquito Larvicide is NOT recommended where the water surface is obscured by vegetation.
	Application: SHAKE WELL BEFORE USING. BIOPREN 50 Liquid Mosquito Larvicide may separate on standing and must be thoroughly agitated prior to dilution. Where possible, use the prepared mixture on the day of mixing.
	1. Ground: Determine the average total spray volume applied per hectare by individual operators and/or specific equipment. Mix BIOPREN 50 Liquid Mosquito Larvicide in sufficient water to deliver the rate per hectare of BIOPREN 50 recommended on this label (220-360 mL/ha).
	2. Aerial: Low Volume (5-50 L/ha) Boom and nozzle - Use the recommended amount of BIOPREN 50 Liquid Mosquito Larvicide in sufficient water to give good coverage. 5-50 L/ha of total spray volume is usually satisfactory. DO NOT apply when weather conditions favour drift from areas treated. A large droplet size (> 150 microns) is recommended to minimise drift and ensure good canopy penetration.
	3. Aerial: Ultra-low Volume (< 5 L/ha) Rotary atomiser - Use the recommended amount of BIOPREN 50 Liquid Mosquito Larvicide in sufficient water to give good atomisation and coverage. Dilution of 1 part BIOPREN 50 to 2 parts water is the recommended minimum. This results in a total application volume of 660 mL (at the 220 mL/ha BIOPREN 50 rate) to 1080 mL (at the 360 mL/ha rate). The quantity of water may be increased to give a total spray volume of up to 5 L to give better coverage. A large droplet size (>150 microns) is recommended to minimise drift and ensure good canopy penetration.
	Mixing: DO NOT mix with oil. Always use clean equipment. Partially fill spray tank with water, then add the recommended amount of BIOPREN 50 Liquid Mosquito Larvicide, agitate and complete filling. Mild agitation during application is desirable.

Resistance Warning:	INSECTICIDE RESISTANCE WARNING GROUP 7A INSECTICIDE For insecticide resistance management Biopren 50 Liquid Mosquito Larvicide is a Group 7A insecticide. Some naturally occurring insect biotypes resistant to Biopren 50 and other Group 7A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Biopren 50 or other Group 7A insecticides are used repeatedly. The effectiveness of Biopren 50 on
	other Group 7A insecticides are used repeatedly. The effectiveness of Biopren 50 on resistant individuals could be significantly reduced. Since occurrence of resistant individuals

	is difficult to detect prior to use, Bábolna Bio accepts no liability for any losses that may result from the failure of Biopren 50 Liquid Mosquito Larvicide to control resistant insects. Biopren 50 Liquid Mosquito Larvicide may be subject to specific resistance management strategies. For further information contact your local supplier, Globe Australia Pty Ltd representative or local agricultural department agronomist.
--	--

Precautions:	Precautions:	
--------------	--------------	--

Protections:	PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS
	DO NOT apply when weather conditions favour drift from treated areas.

Storage and Disposal:	STORAGE AND DISPOSAL KEEP OUT OF REACH OF CHILDREN. Store in the closed, original container in a dry, cool, well ventilated area out of direct sunlight. Store above 5°C. Do not freeze. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the
	surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Safety Directions:	SAFETY DIRECTIONS : May irritate the eyes and skin. Avoid contact with eyes and skin. When preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.
--------------------	---

Australia 131 126.	First Aid Instructions:	FIRST AID : If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.
--------------------	-------------------------	---

First Aid Warnings:	

DIRECTIONS FOR USE

Restraint:

DO NOT apply to dry sites. Biopren 50 Liquid Mosquito Larvicide must be applied directly to water.

SITE	PEST	RATE	CRITICAL COMMENTS
Freshwater swamps and marshes, salt and fresh water lagoons, salt marshes, mangrove	Aedes, Anopheles, Culex and Ochlerotatus	220 mL/ha	Use this rate where water is shallow (< 30 cm), clean, and where larval mosquito counts are low (< 10 per dip).
swamps, estuarine areas, woodland pools, any natural water holding features.	spp.	290 mL/ha	Use this rate where water is deep (> 30 cm), is rich in organic matter and/or sediment, and where larval mosquito counts are high (> 10 per dip).
		360 mL/ha	Use this rate when the major target mosquito species is <i>Culex sitiens</i> .

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION