

READ SAFETY DIRECTIONS BEFORE OPENING  
OR USING



# CHAINDRITE PERMFORCE Dust Insecticide



ACTIVE CONTIITUENT: 10 g/kg PERMETHRIN 25:75

**GROUP 3A INSECTICIDE**

For the control of Cockroaches, Ants, Fleas, Termites, Bedbugs, European Wasps, Spiders  
and other insects in certain situations as per the Directions of Use.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE**

Contents: 2- 15 kg

**APVMA APPROVAL NO: 62664/52573**

**Distributed By:**

Sherwood Chemicals Australia Pty Ltd  
Level 3, 1060 Hay St  
WEST PERTH 6005  
AUSTRALIA  
TEL: 08 9219 4683  
FAX: 08 9219 4672

“Sherwood Logo”

## GENERAL INSTRUCTIONS

CHAINDRITE PERMFORCE dust Insecticide is ready to use, no further diluent be added. Leave powder undisturbed for as long as possible.

## INSECTICIDE RESISTANCE WARNING

<b>GROUP</b>	<b>3A</b>	<b>INSECTICIDE</b>
--------------	-----------	--------------------

For insecticide resistance management CHAINDRITE PERMFORCE Dust Insecticide is a group 3A insecticide. Some naturally occurring biotypes resistant individuals can eventually dominate the insect population if CHAINDRITE PERMFORCE Dust Insecticide or other group 3A insecticides are used repeatedly. The effectiveness of CHAINDRITE PERMFORCE Dust Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Sherwood Chemicals accepts no liability for any losses that may result from the failure of CHAINDRITE PERMFORCE Dust Insecticide may be subject to specific resistance management strategies. For further information, contact your local supplier, Sherwood Chemicals representatives or local agricultural department agronomist.

## PRECAUTIONS

AVOID CONTACT WITH FOOD.

Cover all foodstuffs before application.

**DO NOT** allow dusting powder to contact food, food preparation surfaces or food preparation utensils.

**DO NOT** apply dust directly on humans, exposed food or food utensils.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

Toxic to bees and reptiles. Harmful to fish.

**DO NOT** contaminate dams, rivers, streams, waterways or drains with the product or the used container.

Remove or cover fish tanks before use.

## STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN

Store in the closed, original container in a safe well-ventilated area, as cool as possible. **DO NOT** store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Dispose of rinsings in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots.

If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

## SAFETY DIRECTIONS

Will irritate the eyes, avoid contact with eyes, do not inhale dust. When opening the container and using the product, wear rubber gloves and specific dust, mist or fume face mask covering mouth and nose. Wash hands after use and after each day's use, wash gloves and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). (New Zealand 03 4747000).

## MATERIAL SAFETY DATA SHEET

Additional Information is listed in the Material Safety Data Sheet (MSDS). A Material Safety Data Sheet for CHAINDRITE PERMFORCE Dust Insecticide is available from Sherwood Chemicals on request: 089219 4683

## EXCLUSION OF LIABILITY

Sherwood Chemicals warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with Directions for Use under normal conditions of use. No warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions or under off-label permits not endorsed by Sherwood Chemicals, or under abnormal conditions.

In a Transport Emergency <b>DIAL 000</b> Police or Fire Brigade
---

**APVMA APPROVAL NO: 62664/52573**

**DATE OF MANUFACTURE:**

**BATCH NO:**

## DIRECTIONS FOR USE

Situation		Target Insects	Rate	Critical Comment
Domestic, Institutional and Industrial Premises	Food preparation & Storage areas	Cockroaches & Silverfish	10 g/m <sup>2</sup> Applied with a powder blower	Blow powder into all cracks and crevices. Pay particular attention to warm, dark corners around cookers, fireplaces, radiators, stores, pipes, rubbish chutes and places where liquid sprays may not be used.
		Ants		Sprinkle powder along runs, on and around nests. If ants are invading domestic premises apply powder to the routes by which they enter. Repeat as necessary.
	Living & Sleeping areas	Fleas and Carpet Beetles		Dust powder over floor, brushing into carpet pile and blowing into areas where flea larvae develop.
		Bedbugs		Apply powder to bedsteads, bedsprings, mattresses, floor coverings, upholstered furniture, cracks in walls, behind torn wallpaper, joins in woodwork and other harbourages.
Roof voids, wall and floor voids, crawl spaces		Cockroaches, Silverfish, Ants, Fleas, Spiders and bird mites ( <i>Ornithonyssus spp</i> )		Blow dust into all void areas, ensuring all surfaces are thoroughly treated. Repeat application if re-infestation occurs. Remove all birds nests and introduce physical bird barriers to prevent reinfestation of bird mites.
Skin and Hide Preparation		Hide Beetles ( <i>Demestes spp.</i> )		<p>a) Place skins on drying racks and apply dust using a manual or motorized duster.</p> <p>b) Dust exposed edges of dried, baled skins.</p>
Mushroom Beds		Mushroom, Sciariid flies		Dust mushroom beds no more than twice a week to control adult and larval flies. Discontinue dusting when mushrooms reach the size of a 10 cent piece.
Seed Potatoes		Potato Moths ( <i>Phthorimaea operculella</i> )	600 g/tonne or 30g/50kg bag of seed potatoes	<p>Dust thoroughly immediately after harvest while filling storage containers.</p> <p>Treated seed potatoes are to be used only sowing; does not use for food of feed purposes.</p> <p>Treated seed potatoes are to be stored apart from other potatoes and bags or containers should be clearly marked to identify that the contents have been treated.</p>
Large Wasp Nests		European Wasps	125-250 g per nest	Use hand duster and apply powder through entrance holes (125-250g/nest dependent on size of nest and number of entrance holes)
Bee nests in wall cavities		Feral honey bee		Treat at night wherever possible. Operators should wear clothing protective against the wasp including long – sleeved overalls buttoned at wrist and throat, gloves, hat and bee – veil.
Tree stumps in urban areas		Subterranean Termites ( <i>Coptotermes spp.</i> )	250 g per tree	Upon locating nest in tree stump, drill 3 holes with a maximum diameter of 12mm. Apply dust directly to the nest using a carbon dioxide gas cylinder and a trigger gun. After treatment holes should be coated carefully with a fungicide, e.g. copper naphthenate, and sealed with an appropriate chalking compound.

**NOT TO BE USED FOR ANY PURPOSE, OR ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

### WITHHOLDING PERIOD

Mushrooms: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION